



**FIXED OR REMOVABLE
PROSTHESIS ON IMPLANT
INFORMED CONSENT
CERTIFICATE OF CONSENT**

HD.RB.58 YT:14.01.2015 REV.NO:01 REV.T: 18.04.2022 S.NO: 2/2

Dear:(Patient Name Surname):.....(Date of Birth):.....(TC No):.....**BARKODY**

This form asks for your medical and dental history and some personal information. You have the right to be informed about the treatment and procedures to be carried out. The form also contains information about the procedures to be performed in the clinics where you will be treated. The purpose of these explanations is to inform you and ensure your participation in the treatment process in order to improve and protect your oral dental health. After you have learnt the benefits and possible risks of the treatment and procedures, it is up to your decision to consent to the procedure.

Name and Surname: Email Address:
Tel: Gender:

Contact person in case of emergency:

The patient's legal representative:

(Legal Representative: Guardian for those under guardianship, parents or 1st degree legal heirs for minors. Please indicate the degree of closeness next to the name of the patient's relative).

Name Surname:..... Tel No:.....

I n o r d e r to inform you and to obtain your consent to start your treatment, you m u s t read this form, fill in the sections at the end and sign it. Thank you for your participation and your time.

RECOMMENDED TREATMENT



TOOTH TO BE TREATED	DIAGNOSIS	PLANNED TREATMENT
55 54 53 52 51 18 17 16 15 14 13 12 11	61 62 63 64 65 21 22 23 24 25 26 27 28	
48 47 46 45 44 43 42 41 85 84 83 82 81	31 32 33 34 35 36 37 38 71 72 73 74 75	

Implant Prosthesis

To be implemented

Implant prosthesis is an alternative treatment method for traditional dentures. Basically, it r e f e r s t o the placement of specially produced titanium screws in the jaw bones with one or more missing teeth and the attachment of appropriate crown-bridge or removable prostheses to the retaining mechanisms on them. As can be seen, implant prosthesis treatment is a two-step treatment method that requires direct participation of the patient in terms of cost. A good treatment planning is vital for successful treatment and patient satisfaction. The patient's expectations, general health status and the condition of the jaw bones are factors that directly affect the treatment. For this reason, the physician who will perform surgical and prosthetic procedures will examine you together and will want to perform various procedures for planning. Only at the end of this stage can the treatment planning be shared with you and submitted for your approval. Despite all these precautions, it should be kept in mind that biological tissues are being studied and that changes in treatment planning may be necessary due to unforeseen factors at various stages. Before starting the treatment planning, it is important for both the patient's own safety and the physician that patients share their systemic diseases (heart, diabetes, blood pressure, etc.), infectious diseases (such as hepatitis), and medications, if any, with the physician. If the patient is allergic to any substance (penicillin, etc.), he/she should tell before starting treatment. Since these diseases can significantly increase the risk of treatment failure, it is very important to know them in advance. Habits that may cause trauma to teeth/implants such as smoking, alcohol addiction, grinding teeth pose a risk to the success of implant treatments, so be sure to consult your physician about these issues before starting treatment. You may be asked to have various radiological examinations (periapical radiography, panoramic radiography, intraoral tomography) before, during and after treatment for surgical or prosthetic reasons.

TREATMENT AND PROCEDURES:

LOCAL ANAESTHESIA Local anaesthesia is applied to provide pain control during treatments. Allergic reactions, loss of sensation, bleeding, temporary muscle spasms, temporary facial paralysis, emphysema (swelling of the face) may be seen in the patient after local anaesthesia application, although rare. During the effect period of local anaesthesia, eating and drinking is not recommended until the numbness disappears to prevent the formation of wounds in the cheek and lip due to biting.

IMPLANT TREATMENT: Implants made of titanium similar to the root form are placed in the jaw bones and prostheses are planned to replace the lost teeth. The aim of this treatment is to treat the problem caused by tooth loss and to increase chewing efficiency. Implants are placed surgically. An incision is first made on the soft tissue in the area where the implant will be placed. The bone is exposed and the implant socket is prepared with special tips. The implant screw is placed in this slot opened in the jawbone, the area is closed with sutures. The stitches are removed after one week, and it is normal to experience mild pain and swelling during this one-week period. It is usually left to heal for 2-4 months, longer if necessary. Parts that will support the prosthesis are attached. The next stage is prosthesis construction.

PROSTHETIC TREATMENT Prosthetic treatment begins at the planning stage of the implant prosthesis treatment. Your expectations for the final prosthesis, your systemic and oral condition and the fact that a significant part of the treatment materials will be paid for by you will influence the prosthetic treatment plan. In the initial phase of treatment, the fees you pay for the purchase of implant screws and surgical procedures are not included in the prosthesis construction fees. You have to pay a separate fee for the fixed or removable prostheses you will have made on your implants. Not all of these prostheses are supported by the Social Security Institution. In this case, the treatment fees will be covered by you.

Especially in cases of complete or partial loss of teeth, it may be necessary to make some prostheses for the surgical phase (surgical templates or some prostheses that you need to use immediately after the surgical phase). Not all of these prostheses are supported by the Social Security Institution. In this case, you will have to pay for the treatment yourself. In cases of partial or total edentulism, you may need to use temporary prostheses following the placement of implants. You may not be able to use these prostheses after the end of treatment. The treatment costs for these prostheses will be covered by you. The construction of partial and total prostheses or fixed prostheses can only be started 10-15 days after the implant healing caps are attached. Some problems in the surgical phase or in the healing process may require a deviation from the treatment plan recommended to you or the use of extra materials to realise this treatment plan. The new treatment plan will be realised with your approval. However, you may still have to pay again for some materials to be used in this planning. The end of the prosthetic treatment does not mean that you will not come to our clinic again. The control appointments that will be notified to you by our physician are important. Daily care and home care are important for implant prostheses. In order to use prostheses and implants for many years, it is vital for the use of prostheses that you come to control appointments and apply cleaning in accordance with your physician's recommendations. All prostheses show fatigue in various ways over time. Depending on this fatigue, some parts of your prosthesis may need to be renewed and replaced over time. As with traditional prostheses, implant prostheses may also need to be renewed over time. As long as the implants are not damaged, this does not require any procedure other than the replacement of the prosthesis. However, you may have to pay a fee for these treatments.

Alternative Treatments:Your missing teeth can be treated with classical methods other than implant prostheses. Alternative treatments to implant-supported prostheses are fixed or removable prostheses. The most suitable of these prostheses varies from person to person depending on factors such as the number of remaining teeth, which teeth the remaining teeth are, their location in the mouth, their placement, the health of your teeth, the importance you attach to your oral care, your general health status. Ask your doctor for information about the prosthesis options suitable for you and the advantages and disadvantages of these options.

Expectations from the treatment: The purpose of placing dental implants for the purpose of making fixed or removable prosthesis on the implant is to provide support and retention to crowns, bridges, removable or fixed prostheses i n the absence of natural teeth.

Patient/Guardian



**FIXED OR REMOVABLE
PROSTHESIS ON IMPLANT
INFORMED CONSENT
CERTIFICATE OF CONSENT**

HD.RB.58	YT:14.01.2015	REV.NO:01	REV.T: 18.04.2022	S.NO: 2/2
----------	---------------	-----------	-------------------	-----------

Potential Problems and Risks

Many problems may occur in implant fixed and removable prostheses that vary from patient to patient. Some of these problems are listed below:
Infection in the gingiva around the implant, bone loss in the neck area of the implant, bone loss around the implant, partial or complete loss of the connection of the implant with the bone, necessity of surgical removal of the implant due to this

- Screw loosening, screw jamming, aspiration due to loosening of screws, swallowing
- Screw breakage
- Fractures in the neck area of the implant
- Fractures in the upper part of the implant called abutment
- Complete removal of the implant as a result of its inability to connect with the bone

Fixed implant prostheses may cause many different problems that are not limited to the problems mentioned below but vary from patient to patient, as well as problems similar to prostheses applied to natural teeth (see consent form for fixed prostheses)

- In cases such as insufficient amount of bone, inability to achieve the ideal aesthetic appearance and inability to provide adequate support to the cheek and lip, appearance of the neck area of the implant, metal colour from the gum, restrictions that may occur in relation to the colour or shape of the prosthetic teeth.
- Dislocation of the capping material, screw loosening, dislocation, loss in screw crown and bridge prostheses.
- The prosthesis cannot be positioned appropriately as a result of the implant not being ideally positioned or the implant cannot be used as a support for the prosthesis.
- Food entrapment, air leakage in the areas prepared for cleaning by the patient in implant prostheses
- Loosening or dislocation of the crown or bridge on the implant supports
- Fractures and breakages in crown or bridge materials (porcelain, metal, etc.)
- Occlusal adaptation may be required for prostheses
- Need for replacement of crowns and bridges

In removable implant prostheses, besides many different problems that are not limited to the problems mentioned below but vary from patient to patient, problems similar to prostheses applied to natural teeth may also occur (see consent form for removable and total prostheses):

- Food accumulation under the prosthesis
- Prostheses cause dents
- Wear and deterioration of the implant retainers, rubber or metal parts due to use, deterioration of the retainer parts due to improper use and the need for replacement. Replacement of these parts is mandatory and requires the patient to pay an additional fee.
- Dislocation or loss of the attachment part of the prosthesis. In this case, replacement is mandatory and the patient has to pay an additional fee.
- Decrease in prosthesis retention over time, loosening, need for feeding in prostheses.
- Fractures or cracks in prostheses
- Problems with speech
- Occlusal alignment needs
- Fractures or cracks in the dentures, protrusion or fracture of the denture teeth, wear over time.
- Colouration of the prosthesis material, deterioration or aging over time or due to improper use, the need for complete replacement of the prosthesis.

In case of any problems that may occur with implant prostheses, you should definitely consult your doctor as soon as possible.

If the recommended treatment is not applied: In cases where restoration of tooth deficiencies with implant prostheses cannot be performed, it is recommended that you have your tooth deficiencies treated with other classical approaches that your doctor deems appropriate.

After the completion of the implant prosthesis, you should consult your doctor for control at regular intervals such as every 6 months-1 year or at regular intervals recommended by your doctor. Always consult your doctor about how to maintain the hygiene of your prostheses and implants. It is the sole responsibility of the patient to fulfil your doctor's recommendations completely, to protect and maintain the hygiene of your implants, prostheses and natural teeth and gums, to come to routine control examinations, and to report your problems as soon as possible. Habits that may cause trauma to teeth / implants such as smoking, alcohol use, teeth grinding, excessively hard and sticky foods negatively affect the success of the implant and prosthesis. After the completion of your prosthesis, it is entirely your responsibility to avoid these situations, to fulfil your doctor's recommendations on related issues and the problems that may occur if you do not bring them.

Special Cases: You can reach Süleyman Demirel University Faculty of Dentistry doctors by phone on 0246 211 3352 or by visiting.

APPROVAL

My whole mouth was examined in detail. In addition, in the relevant sections, the physicians explained what oral dental disease is, why treatment is needed, the risks involved, the problems that may occur, alternative methods, changes that may occur after treatment, the possibility of success and the situations that may occur during the healing process.

During the dentist's diagnosis and treatment;

- ✓ Consultation may be requested from other physicians and they may be involved in the treatment process,
- ✓ Assistants and trainee dentists will participate in diagnosis and treatment, and trainee dentists will work under the supervision of assistants and faculty members,
- ✓ During the diagnostic method, intervention or treatment, local anaesthesia may be applied as part of these procedures, attendance to the appointments given without delay and compliance with the physician's recommendations and practices regarding the treatment may directly affect the treatment results,
- ✓ Oral, Dental and Maxillofacial Radiology doctors, trainee dentists, dental technicians and x-ray technicians can take films,
- ✓ My anamnesis information and radiological images, photographs, examination results (pathology report, laboratory results, etc.) can be used for diagnostic, scientific, educational or research purposes by keeping my identity information confidential,

It was explained to me.
I was informed about the Implementation Communiqué published by the Social Security Institution (SSI) for the relevant treatments. I was informed that the invoices for the items that the SSI does not pay for the material to be used during the stages of the treatment should be paid by me

Dear Patient, You have the right to be informed about your health condition and the benefits/harms and alternatives of the proposed diagnostic or therapeutic procedures; to accept or partially/completely reject the treatment; and to stop the procedures at any stage! This document, which we want you to read and understand, has been prepared not to frighten you or to keep you away from medical practices, but to inform you, to determine whether you consent to these practices and to obtain your consent. This consent form consists of 2 pages and has been issued in 2 copies in accordance with Article 70 of the Law No. 1219 and Article 26 of the Turkish Penal Code No. 5237, and one copy has been given to the patient / patient's legal representative.

I HAVE BEEN TOLD ABOUT THE TREATMENT FOR MY DISEASE AND I HAVE READ AND UNDERSTOOD THIS FORM. I ACCEPT THE TREATMENT (Note:

Please write in **YOUR OWN HANDWRITING** in the space below).

PATIENT'S/RELATIVE'S/GUARDIAN'S

FULL NAME :.....

DATE:

TIME:.....

SIGNATURE

THE PHYSICIAN PROVIDING THE INFORMATION

FULL NAME.....

DATE:.....

CLOCK:.....

SIGNATURE/SIGNATURE